From Aerosols to Populations

Jonathan M. Samet, MD, MS

Dean and Professor

Colorado School of Public Health


EMHI Airborne Transmission of SARS-CoV-2
Virtual Workshop

August 26-27, 2020

colorado school of public health

UNIVERSITY OF COLORADO
COLORADO STATE UNIVERSITY
UNIVERSITY OF NORTHERN COLORADO

Bringing the story together



Putting H-D together



Population Measures

- Incidence (count)
- Test positive
- Hospitalization
 - ICU need
- Mortality
- R_0 and R_e

Relationship to airborne transmission?

R_o: The basic reproductive number

Definition: the average number of secondary cases generated by a primary case in a 100% susceptible population.

$$R_0 = CPD$$


C=number of contacts per unit time
P=probability of infection when a
contact occurs
D=duration of infection

Comparing SARS-CoV-2 with SARS-CoV and influenza pandemics


Eskild Petersen, Marion Koopmans, Unyeong Go, Davidson H Hamer, Nicola Petrosillo, Francesco Castelli, Merete Storgaard, Sulien Al Khalili, Lone Simonsen

	SARS-CoV-2	SARS-CoV	Pandemic influenza 1918	Pandemic influenza 2009	Interpretation
Transmissibility, R₀	2.5	2.4	2.0	1.7	SARS-CoV-2 has the highest average <i>R</i> ₀
Incubation period, days	4–12	2–7	Unknown	2	Longer incubation period; SARS-CoV epidemics form slower


Systematic Review: Measles R₀ Estimates by Geographical Location and Time



The Effective Reproduction Number Estimates Based on Laboratory-Confirmed Coronavirus Disease...




Colorado: the estimated reproductive number from March to now



This is based on estimation method 2 in the table on the next page

R_e: Effective reproductive number


Definition: Describes transmission once the epidemic is established. **Population**




R_e is a Multi-Level Parameter

Macro-environment


Microenvironment ersonal



The SEIR Model



The SEIR Model Elaborated



The SEIR Model and CQ1-4

